

auditory system:

- very fast processing (visual system slower)

↳ 2nd messengers etc.

- very sensitive: process 2 components of sound

1. amplitude of sound wave (decibels)

- human ear detects $0 \rightarrow 120$ dB, can detect 0.01 nm shift in air

frequency



amplitude


2. frequency of sound (Hz)

- $20 - 20,000\text{ Hz}$: speech $\sim 1000 - 3000\text{ Hz}$

- $50\mu\text{sec}$ detection (faster than synaptic transmission?) $\leftarrow 1\text{ ms}$


- more complex than visual system

in cochlea:



hair cell
(like photoreceptor
in visual system).

in brainstem:



ganglion cell

in thalamus:


(sound loc.)
Superior olivary
nucleus

in thalamus:


medial
geniculate
nucleus

1^{o} auditory
cortex
(Brodmann's
area 41/42):

topological
by frequency
in superior
temporal gyrus



inferior
collicular
(also helps
determine where
sound coming from
(in midbrain)



everything ipsi- & contralateral
past cochlear nucleus

(to get hearing loss in one ear,
must affect path before cochlear
nucleus)

past this, must have bilateral
lesions for hearing loss

contralateral projections
(each to other auditory
olivary nucleus), allows
to compare inputs from
each ear



association area


Wernicke's area: important for language,
unilateral lesion (esp. on left) can have
devastating effects

- hair cells have graded potentials (like PRs & bipolar cells)

- ganglion cells fire APs

(2 muscles that can reduce sound?)
air-filled

semicircular canals (balance)



- directs sound towards ear canal

(damage still lets you localize sound b/c one ear can do this somewhat)

if get ear infections, can get middle ear scar tissue


(common in children)

otosclerosis in old people, bone spurs in connective tissues
(can connect w/ prosthetic stapes)

or damage to cochlea, certain frequencies you hear

unfold cochlea:


over & over



Specific frequency will give vibration at specific area


- hair cells have place code

cochlear cross-section:



- as basilar membrane goes up, //

kinocilium goes down, //



ganglion cell

primary auditory nerve (can record from this)

(activity)

no sign change:

beta fire at slow constant rate;
more w/ sound,
less when sound off

unusual b/c charge carried by only one ion (Cl^-)


- this helps make it fast:

- direct gating of ion channel (cur timescale)

- K^+ can come in & leave

ion channel that mediates mechanotransduction yet to be cloned in humans

- trp channel
(K^+ channel)




S4 voltage sensor (not in trp)

trp channels - 6 TM

- no S4

- very unusual N-termini (long, w/ ankyrin repeats)


↳ motif that binds cytoskeleton (eg actin)




- nonselective cation channel

- when hair bundle shifted, little bit of Ca^{2+} goes in channel (along w/ K^+)

- adaptation to constant sound




- change in way channel is gated



during prolonged Ca^{2+} entry,
binds myosin, drags down actin
cable, channels start to close:

now tip link doesn't activate channel anymore




stimulus stimulus locking:

hair cell V_m oscillates w/
movement of stereocilia

($\rightarrow 3 \text{ kHz}$)

in upper range, doesn't keep up but
talks to many ganglion cells




each hair cell talks to many (10-20) ganglion cells


- need dense body (particular to hair cells): like conveyor belt
to release NT very rapidly

frequency
of firing matches
Stimulus oscillation


place code: depending on where hair cells are, encode particular frequencies

Stereocilia length
also help tune to
specific frequencies






- aminoglycoside antibiotics (eg kanamycin, streptomycin) block mechanosensitive channels, can kill hair cells over time
- loud noises can shear stereocilia: worst if same frequencies over & over
 - correct w/ hearing aids at base of eardrum (tune to defective frequencies)
 - can also use cochlear implants: if ganglion cells still alive



↳ takes place of basilar membrane frequency decoder, excite correct electrodes, stimulate correct axons



7.29J / 9.09J Cellular Neurobiology

Spring 2012

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.